

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



(51) International Patent Classification ⁶ : C12N 15/09		A1	(11) International Publication Number: WO 98/45426
			(43) International Publication Date: 15 October 1998 (15.10.98)
(21) International Application Number: PCT/US98/06897 (22) International Filing Date: 6 April 1998 (06.04.98) (30) Priority Data: 08/833,814 9 April 1997 (09.04.97) US (71) Applicant: ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US). (72) Inventors: HEIDLER, Steven, A.; 9912 Dairying Street, Fishers, IN 46038 (US). RADDING, Jeffrey, A.; 1424 Ironwood Drive West, Carmel, IN 46033 (US). YOUNG, Debra, A.; 1870 Ginseng Trail, Danville, IN 46122 (US). (74) Agents: WEBSTER, Thomas, D. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: SCREEN FOR INHIBITORS OF FUNGAL IPC SYNTHASE (57) Abstract <p>This invention provides novel fungal strains altered in IPC synthase production, and a method for screening for inhibitors of fungal IPC synthase using whole cells.</p>			
ATTORNEY DOCKET NUMBER: 10182-016-999 SERIAL NUMBER: 10/032,585 REFERENCE: BQ			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

SCREEN FOR INHIBITORS OF FUNGAL IPC SYNTHASE

5 BACKGROUND OF THE INVENTION

This invention relates to the analytical arts and to recombinant DNA technology. In particular the invention pertains to the construction and purification of novel strains of yeast and other fungi that are useful in whole-cell screens for inhibitors of inositolphosphoryl ceramide synthase (IPC Synthase).

The incidence of life-threatening fungal infections is increasing at an alarming rate. With the exception of *Staphylococci* infections, the fungus *C. albicans* represents the fastest growing area of concern in hospital acquired infections. About 90% of nosocomial fungal infections are caused by species of *Candida* with the remaining 10% being attributable to *Aspergillus*, *Cryptococcus*, and *Pneumocystis*. While effective antifungal compounds have been developed for *Candida* there is growing concern that the rise in fungal infections may portend a trend toward escalating resistance and virulence in the future. This is problematic because anti-*Candida* compounds rarely possess clinically significant activity against other fungal species.

25 Inositolphosphoryl ceramides are sphingolipids found in a number of fungi including but not limited to *S. cerevisiae*, *S. pombe*, *C. albicans*, *A. fumigatus*, and *H.*

-2-

capsulatum. A step of sphingolipid biosynthesis that is unique to fungi and plants is catalyzed by the enzyme IPC synthase. The IPC synthase step, which covalently links inositol phosphate and ceramide, is essential for viability in *S. cerevisiae*. Although some elements of sphingolipid biosynthesis in fungi are shared with mammalian systems, the pathways diverge after formation of ceramide. Thus, the formation of inositolphosphoryl ceramide is unique to fungi and plants, making IPC synthase a good molecular target for antifungal chemotherapy.

While IPC synthase presents a rational target for antifungal therapy, presently there are no clinically useful compounds that act at this step. Thus, there is a need for new compounds that inhibit IPC synthase.

BRIEF SUMMARY OF THE INVENTION

The present invention relates to fungal IPC synthase and to whole-cell screens for inhibitors thereof.

In one embodiment the invention relates to novel fungal strains in which IPC synthase is produced at levels that differ from wild-type cells.

In another embodiment, the invention relates to novel fungal strains in which a gene that encodes IPC synthase or subunit thereof is operably-linked to a heterologous promoter such that IPC synthase is produced at levels that differ from wild-type cells.

In another embodiment the present invention pertains to high throughput screens for inhibitors of fungal IPC synthase.

DESCRIPTION OF THE DRAWINGS

Figure 1. Plasmid pDAY307 is a 2 μ yeast vector that carries a modified *aur1*⁺ gene. pDAY307 was constructed in vector pYX212 (available from Novagen, 597 Science Dr., Madison, WI 53711), which is a multi-copy vector that carries the URA3 gene and cloning sites for expressing a gene from a TPI promoter.

Figure 2. Plasmid pDAY309 is a 2 μ yeast vector that carries a modified *aur1'* gene. pDAY309 was constructed in Novagen vector pYX213, which is a multi-copy vector that carries the URA3 gene and provides cloning sites for expressing heterologous genes from a GAL1 promoter.

DEFINITIONS

"*aur1'*" refers to a gene of yeast that when mutated confers resistance to aureobasidin A. Aureobasidins are antifungal cyclic peptides isolated from *Aureobasidium pullulans* R106-1 (Takesako et al. *J. Antibiot.* 44, 919-924, 1991). *aur1'* encodes IPC synthase or a subunit of the IPC synthase enzyme. Aureobasidin A (R106-1) inhibits IPC synthase activity *in vitro*.

" Δ *aur1'*" refers to a deletion mutation in the *aur1'* locus.

"Hypersensitive" refers to a phenotype in which cells are more sensitive to antifungal compounds than are wild-type cells of similar or identical genetic background. As used herein, "hypersensitive" means that cells are at least 20-fold more sensitive than wild-type, when grown under identical conditions.

"TPI" refers to the yeast triose phosphate isomerase gene. The TPI promoter is used as a strong constitutive promoter for expression of heterologous genes in yeast (R. Schiestl and R. Gietz, *Curr. Genet.* 16, 339-46, 1989).

"GAL" refers to a structural gene associated with galactose metabolism in yeast. The GAL1 gene, for example, encodes the enzyme galactokinase. The GAL1 promoter, when operably-linked to a heterologous gene, is useful for providing galactose inducibility and glucose repressibility (M. Johnston and R.W. Davis, *Mol. Cell. Biol.* 4, 1440-48, 1984).

"nmt" (no message in thiamine) refers to thiamine responsive yeast promoters of various strengths.

-4-

"YEPD" refers to a rich medium for growing yeast cells. YEPD comprises 1% yeast extract, 2% Bacto-peptone, and 2% glucose.

5 "SC-URA" refers to a synthetic complete medium that lacks uracil. It comprises, per liter, 1.7 g yeast nitrogen base (without amino acids and ammonium sulfate), 5.0 g ammonium sulfate, 0.8 g CSM-URA (available from BIO 101, 1070 Joshua Way, Vista, CA 92083), and 20 g of any suitable carbon source (e.g. glucose, raffinose, sorbitol, or
10 galactose).

The term "deletion mutation" as used herein includes mutations that remove some or all of the DNA that comprises a structural gene or regulatory region for such gene. Also contemplated by the term are mutations that remove DNA from
15 a structural gene and/or its regulatory region and insert therefore a heterologous fragment of DNA that may encode another gene.

"Knockout cassette" means a fragment of native chromosomal DNA having cloned therein a foreign piece of DNA
20 that may provide a selectable marker.

"Recombinant DNA cloning vector" as used herein refers to any autonomously replicating or integrating agent, including, but not limited to, plasmids and phages, comprising a DNA molecule onto which one or more additional
25 DNA segments can or have been added.

The term "expression vector" as used herein refers to any recombinant DNA cloning vector, for example a plasmid or phage, in which a promoter and other regulatory elements are present to enable transcription of the inserted DNA.

30 The term "vector" as used herein refers to a nucleic acid compound used for introducing exogenous DNA into host cells. A vector comprises a nucleotide sequence which may encode one or more protein molecules. Plasmids, cosmids, viruses, and bacteriophages, in the natural state or which
35 have undergone recombinant engineering, are examples of commonly used vectors.

The term "whole-cell screen" refers to the use of whole fungal cells, for example yeast cells, in screens for compounds that inhibit or otherwise alter the growth of said cells on a defined medium. As used herein the term relates
5 to screens for inhibitors of fungal IPC synthase using yeast or other fungi that produce altered levels of IPC synthase or a subunit of the IPC synthase enzyme.

DETAILED DESCRIPTION OF THE INVENTION

10 In yeast and other fungi, such as *C. albicans*, *A. fumigatus*, *C. neoformans*, *C. krusei*, *C. parapsilosis*, *C. tropicalis*, and *C. glabrata*, for example, IPC synthase catalyzes a step in the synthesis of inositolphosphoryl ceramide from ceramide and phosphatidylinositol (G. Becker
15 and R. Lester, Biosynthesis of phosphoinositol-containing sphingolipid from phosphatidylinositol by a membrane preparation from *Saccharomyces cerevisiae*. *J. Bacteriol.* 142, 747-754, 1980). Sphingolipids are necessary for growth and viability of the yeast *S. cerevisiae*. Since IPC synthase is
20 unique to fungi and plants it is a good target for antifungal therapy.

This invention relates to the use of fungal cells in whole-cell screens for inhibitors of IPC synthase. The methods contemplated herein relate to cells that have been
25 modified to express more or less IPC synthase activity than wild-type cells. Cells that are suitable for this purpose could arise, for example, through alterations in the expression of the gene that encodes IPC synthase or subunit thereof, or through alterations in the IPC synthase enzyme
30 itself. Cells having these characteristics can be generated through chemical or UV mutagenesis by well known means, or by recombinant DNA techniques in which, for example, an IPC synthase gene is operably-linked to a heterologous promoter. Transcription from a heterologous promoter is expected to
35 alter expression of an operably-linked gene. The methods contemplated herein are preferably carried out by recombinant DNA techniques.

-6-

The genes that encode IPC synthase or subunit thereof from *S. cerevisiae*, *S. pombe*, and *C. albicans* have been described (EP O 644 262 A2). Methods for introducing mutations into cloned DNA and for linking genes to
5 heterologous promoters are well known to the skilled artisan (See generally, Maniatis et al. Molecular Cloning, Cold Spring Harbor, 1982). A number of suitable heterologous promoters are contemplated including the TPI promoter, the GAL promoters, and nmt promoters, for example. The skilled
10 artisan also recognizes that methods and techniques other than those exemplified herein could be employed to accomplish the same objective and these are intended to be within the scope of the present invention.

The methods disclosed herein relate to the use of whole
15 cells that express altered levels of IPC synthase. For example, cells that express less than wild-type levels of IPC synthase may become more sensitive to inhibitors of IPC synthase. In a preferred embodiment, the methods disclosed herein use cells that are hypersensitive to IPC synthase
20 inhibitors such as R106-1. Hypersensitivity may be achieved, for example, by introducing into fungal cells an expression vector that carries the gene encoding IPC synthase or subunit thereof, such that expression of said gene can be regulated, or maintained at levels that differ from wild-
25 type. The skilled artisan will recognize a number of suitable expression vectors that incorporate a variety of ARS, 2 μ , or CEN replication and segregation elements for expression in *S. cerevisiae* (See e.g. R. Sikorski & P. Hieter, *Genetics*, 122, 19-26, 1989).

30 The present invention contemplates application to any suitable fungal cell. Examples of suitable cells include *C. albicans*, *A. fumigatus*, *C. neoformans*, *C. krusei*, *C. parapsilosis*, *C. tropicalis*, *C. glabrata*, *S. cerevisiae*, and *S. pombe*. Recombinant DNA techniques such as transformation
35 and the like are available for these cells (See e.g. J. Pla et.al. *Yeast*, 12, 1677-1702, 1996; S. Moreno et.al. *Meth. Enzymol.* 194, 795-823, 1991). Without intending to limit the

-7-

scope of the invention, the method will be described in detail with reference to *S. cerevisiae*.

In *S. cerevisiae*, the *aur1*⁺ gene encodes IPC synthase or a subunit of IPC synthase. The *aur1*⁺ recombinant strains
5 described in this invention were created in an Δ *aur1*⁺ genetic background. Since the Δ *aur1*⁺ genotype is lethal in haploid cells on all media tested, it was necessary to begin with a diploid strain in order to create an Δ *aur1*⁺ deletion at one of the chromosomal alleles. Strain YPH501 was used
10 for this purpose (MATa/ α ura3-52 lys2-801 ade2-101 trp1- Δ 63 his3- Δ 200 leu2- Δ 1; R. Sikorski & P. Hieter, *Genetics*, 122, 19-26, 1989).

An Δ *aur1*⁺/*aur1*⁺ heterozygous diploid was produced by transforming YPH501 with a knockout cassette comprising
15 three elements: at the two ends of the cassette are sequences that flank *aur1*⁺ on the yeast chromosome; in the center of the cassette is any suitable marker, for example the TRP1 gene. Other genetic marker genes, well known to the skilled artisan, could also be used for this purpose, for
20 example, HIS3, URA3, and LEU2. Using the knockout cassette, Trp⁺ transformants of YPH501 were selected and tested by Southern hybridization to identify isolates that had undergone a recombination event at the *aur1*⁺ locus.

A suitable heterozygous diploid having the desired
25 *aur1*⁺::TRP1/*aur1*⁺ genotype was identified by sporulation and tetrad analysis, revealing 2 viable spores and 2 nonviable spores. This isolate was transformed with a 2 μ yeast vector (either pDAY307 or pDAY309) that carries a modified *aur1*⁺ gene. The pDAY vectors were constructed in Novagen vectors
30 pYX212 and pYX213, which are multi-copy vectors carrying the URA3 gene and cloning sites that enable expression from TPI(pDAY307) or GAL1(pDAY309) promoters (Novagen, Inc. 597 Science Drive, Madison, WI 53711). The protein that is produced by the *aur1*⁺ gene on the pDAY vectors is modified
35 by the addition of 14 amino acids at the N terminal side of the initiator Methionine. Additionally, the *aur1*⁺ gene on the pDAY vectors is operably-linked to the TPI promoter

-8-

(pDAY307) or to the GAL1 promoter (pDAY309). The *aur1*⁺ gene can be isolated most conveniently by the polymerase chain reaction (PCR) using suitable oligonucleotide primers. The nucleotide sequence for *aur1*⁺ is available from The Stanford
 5 Genome World Wide Web site (URL:<http://genome-www.stanford.edu/Saccharomyces/>); the sequence of an R106-1 resistant allele is also available, See e.g. Heidler and Radding, *Antimicrob. Agents Chemother.* 39, 2765-69, (1995). The pDAY vectors enable regulated expression of a functional
 10 IPC synthase as evidenced by rescue of *aur1*⁺::*TRP1* cells from nonviability. Suitable strains were isolated by sporulation, dissection, and identification of *aur1*⁺::*TRP1*pDAY307 or pDAY309-containing Ura⁺ segregants.

The *S. cerevisiae* strains described in this invention
 15 are listed in Table 1 along with their relevant genotypes.

Table 1

<u><i>aur1</i>⁺-Modified</u>		
<i>S. cerevisiae</i>	Strains	Genotype
20 YPH499Δ <i>aur1</i> ⁺ (pDAY307)	Strain A	MATa <i>ura3-52 lys2-801 ade2-101</i> <i>trp1-Δ63 his3-Δ200 leu2-Δ1</i> <i>aur1</i> ⁺ :: <i>TRP1</i> pDAY307
25 YPH499Δ <i>aur1</i> ⁺ (pDAY309)	Strain B	MATa <i>ura3-52 lys2-801 ade2-101</i> <i>trp1-Δ63 his3-Δ200 leu2-Δ1</i> <i>aur1</i> ⁺ :: <i>TRP1</i> pDAY309

30 The yeast strains relating to the present invention afford regulatable expression of the *aur1*⁺ gene in *S. cerevisiae*. Strain A, for example, provides expression of a vector-borne *aur1*⁺ gene operably-linked to the TPI yeast promoter. This construct results in constitutive expression
 35 of IPC synthase.

Strain B, on the other hand, provides a vector-borne *aur1*⁺ gene whose expression is regulatable. The plasmid-

-9-

borne *aur1*⁺ gene in Strain B is operably-linked to the GAL1 yeast promoter, which imparts inducibility when cells are grown on galactose; moreover, the GAL1 promoter imparts glucose repressibility *in vivo* leading to lowered expression of IPC synthase.

High-Throughput Screen for IPC Synthase Inhibitors Using *aur1*⁺- Altered Fungal Strains

The strains contemplated by the present invention are useful in whole-cell screens for compounds that inhibit fungal IPC synthase. In one embodiment, test compounds are assayed for their affect on the rate of growth or viability of suitable cells. In a preferred embodiment of this method wild-type cells are compared against cells that manifest altered IPC synthase activity, via lowered expression of a gene encoding IPC synthase or subunit thereof.

For example, *S. cerevisiae* strains A and B (Table 1) are equally affected by antifungal compounds that do not target IPC synthase, such as amphotericin B, nystatin, echinocandin analog, cycloheximide, fluconazole, and ketoconazole (See Table 3).

In contrast, strains A and B differ markedly in their sensitivity to R106-1, a known inhibitor of IPC synthase. When strain A (which provides expression of *aur1*⁺ from a TPI promoter) is exposed to R106-1, cells exhibit wild-type sensitivity on all media tested (see Table 2). On the other hand, strain B (*aur1*⁺ expression linked to the GAL1 promoter) exhibits hypersensitivity to R106-1 when cells are grown in a medium containing glucose (Table 2). Similar results were obtained using raffinose as the carbon source. Thus, strain B is at least 30-fold more sensitive than wild-type cells to R106-1, a compound known to inhibit IPC synthase.

This invention further contemplates the use of whole cells in high throughput screens of natural products, synthetic compounds, and compounds derived from combinatorial libraries to identify inhibitors of IPC

-10-

synthase. For this purpose, a comparison is made between, for example, wild-type cells and cells in which IPC synthase gene expression is altered, preferably so that said alteration results in hypersensitivity to inhibitors of IPC synthase, such as R106-1. A screening system of this format would comprise the steps of:

- a) cultivating suitable fungal cells on an appropriate liquid or solid medium;
- b) exposing said cells to a test compound;
- 10 c) monitoring growth inhibition by any suitable means; and
- d) comparing the inhibition of growth of wild-type cells with cells in which IPC synthase expression is modified.

15 In a preferred embodiment the cells are derived from strains A and B. This screening system may be adapted to automated procedures such as a PANDEX[®] system (Baxter-Dade Diagnostics), allowing for efficient high-volume screening of potential inhibitors.

20 The following examples more fully describe the present invention. Those skilled in the art will recognize that the particular reagents, equipment, and procedures described are merely illustrative and are not intended to limit the present invention in any manner.

25

EXAMPLE 1

Construction of an *S. cerevisiae* strain exhibiting altered IPC synthase expression

A deletion of the *aur1*⁺ locus of *S. cerevisiae* strain YPH501 was constructed as follows. First, a knockout cassette was constructed in order to replace the wild-type *aur1*⁺ chromosomal locus with the *TRP1* gene. Using oligonucleotide primers, designated SEQ ID NO.1 and SEQ ID NO.2, a 909 base pair fragment of chromosomal DNA flanking the 3' end of *aur1*⁺ was PCR-amplified from yeast genomic DNA. Next, a 1086 base pair fragment was amplified from yeast chromosomal DNA flanking the 5' end of the *aur1*⁺ locus

-11-

using primers designated SEQ ID NO.3 and SEQ ID NO.4. The PCR-amplified genomic fragments were ligated to a *TRP1* gene (St.John et al. *J.Mol.Biol.* 152, 317-34, 1981) to form the knockout cassette used in the next step to delete the *aur1*⁺ locus.

The *TRP1* knockout cassette was transformed into diploid strain YPH501. This strain carries a deletion mutation at the chromosomal *TRP1* locus. Trp⁺ transformants were selected by plating cells onto synthetic complete medium, or minimal medium plus supplements lacking tryptophan. A transformant having the desired gene transplacement at the *aur1*⁺ locus was identified by sporulation and tetrad analysis. This analysis identified a suitable heterozygote as producing 2 viable spores and 2 non-viable spores. The relevant genotype of this heterozygote is *aur1*⁺::*TRP1/aur1*⁺.

In order to isolate a haploid strain carrying the *aur1*⁺::*TRP1* deletion allele, the aforementioned *aur1*⁺::*TRP1/aur1*⁺ heterozygote was transformed with an expression plasmid that carries the *aur1*⁺ gene in operable linkage with a TPI (pDAY307) or GAL1 (pDAY309) promoter (See Figures 1 and 2). Transformants were sporulated and two isolates having the correct phenotype (viz. Trp⁺ Ura⁺) were chosen for further study (designated Strain A (pDAY307) and Strain B (pDAY309) (See Table 1). Hypersensitivity in Strain B was conferred by the plasmid-borne sequence under the control of the specified promoter. This was confirmed by transformation of Strain B with a second plasmid bearing a wild-type *aur1*⁺ sequence under the control of its own promoter. This transformant regained wild-type sensitivity upon loss of the pDAY309 plasmid.

EXAMPLE 2

Construction of a strain of *S. pombe* exhibiting altered IPC synthase expression

A strain of *S. pombe* exhibiting underexpression of *aur1*⁺ is constructed in a manner analagous to Example 1. A gene replacement cassette is constructed as follows. Using

-12-

PCR amplification, *S. pombe* chromosomal sequences are amplified from the 5' end (primers designated herein as SEQ ID NO 5 and SEQ ID NO 6), and 3' end (primers designated herein as SEQ ID NO 7 and SEQ ID NO 8) of the *aur1*⁺ locus.

- 5 The amplified fragments are ligated onto the ends of a selectable marker, for example, the *LEU2* gene using any suitable cloning linkers. The resulting gene disruption cassette may be cloned into any suitable integration vector (i.e. one lacking a *pombe* origin of DNA replication), for
10 example the pUC, pBR, or Bluescript *E.coli* plasmids.

In order to create an *aur1/aur1::LEU2* heterozygote, the *LEU2* disruption cassette described in preceding paragraph is transformed into a diploid strain having the following genotype: *h-/h+ ura4-D18/ura4-D18 leu1-32/leu1-32 ade6-M210/ade6-M216*. These diploid cells are white in color owing to cross-complementation by the *ade6* alleles. Transformants are selected following the lithium acetate method, on minimal medium lacking leucine. (See Okazaki et al. Nuc. Acid Res. 18, 6485-89 (1990); Moreno et al. Meth. Enzym. 194,
15 795-823 (1991).

An expression vector that carries the wild-type *aur1*⁺ gene minus the gene's native promoter is cloned behind an *nmt* promoter, as follows. A 1383 base pair fragment of *pombe* genomic DNA carrying the *aur1*⁺ gene is amplified using the
25 primers designated herein as SEQ ID NO 9 and SEQ ID NO 10. These primers carry *Bam*HI and *Xho*I cloning sites, which when digested with said restriction enzymes, are suitable for cloning into a REP "X" vector. The REP X vector carries the *URA4* gene, an *nmt* promoter, and a polylinker cloning site
30 that lacks an ATG codon. (See e.g. Basi et al. Gene 123, 131-36 (1993); Forsburg, Nuc. Acid Res. 21, 2955-56 (1993); Maundrell, J. Biol. Chem. 265, 10857-864 (1990).

The *aur1*⁺ expression plasmid described above is transformed into the *aur1/aur1::LEU2* heterozygote and
35 transformants are selected on minimal medium lacking uracil, leucine, and thiamine. Transformants selected thusly are induced to sporulate by continued incubation. Haploid

-13-

colonies are screened for by identifying "red" colonies (ade6 mutation leads to red color; segregation of complementing ade6 alleles in the diploid produces red haploid colonies). The haploid isolates are leu⁺ ura⁺ and
5 express the aur1 gene product solely from the plasmid borne nmt promoter.

By growing the haploid constructs in medium that contains increasing amounts of thiamine one can identify a level of thiamine that represses expression of aur1⁺ to the
10 lowest levels possible while retaining viability. These cells are expected to be hypersensitive to IPC synthase inhibitors, such as R106-1, and useful for the whole-cell screens described herein.

15

EXAMPLE 3

Differential Sensitivity of Strains A and B to R106-1

Overnight cultures of Strain A and Strain B were grown in YEPD. An aliquot of each overnight culture was diluted into fresh medium. Then about 10⁴ cells in 100 μ l of medium
20 were deposited into each well of a 96-well microtiter plate containing about 100 μ l per well of medium plus decreasing amounts of R106-1, starting from 2.5 μ g/ml and diminishing by successive 2-fold dilutions (R106-1 is available from Panvera Corp. 545 Science Drive, Madison, WI 53711).
25 Innoculated plates were placed in a 30°C incubator and checked for cell growth at 24 hours and 48 hours. Cells were resuspended after 48 hours and allowed to settle for at least one hour, when the cell density was determined at 590 nm using a microtiter plate reader.

30 The results of this experiment are summarized in Table 2. Strain B is from 30-fold to 100-fold more sensitive to R106-1 than either wild-type cells or cells from Strain A. Similar results were obtained using SC-URA (2% glucose) medium (Table 2), and when the cells were grown in medium
35 containing 2% raffinose rather than 2% glucose (data not shown).

-14-

Table 2

Sensitivity of Strains A and B to R106-1
(Minimum Inhibitory Concentration, ug/ml)

5	Strain	SC-URA Glucose	YEP Glucose
	YPH499 ^a	0.625	0.04
	Strain A	0.625	0.04
	Strain B	0.015	0.0006

10 ^a wild-type strain transformed with YEp352 (a vector that carries URA3 gene but not *aur1*⁺; Hill et al. Yeast, 2, 163-67, 1986).

EXAMPLE 4

15 Whole-Cell Screen of Yeast Strains using Non-IPC Synthase
Inhibitors

Strains A and B, and wild-type strain YPH499 (transformed with YEp352) were tested for sensitivity to a variety of known antifungal compounds. Cells were grown in
20 YEPD or SC-URA (2% glucose) liquid culture and dispensed into the wells of a 96-well microtiter dish containing decreasing amounts of antifungal compounds, as in Example 2.

The results of this experiment demonstrate that strains A and B behave similarly to wild-type cells when exposed to
25 antifungal compounds that do not target IPC synthase (See Table 3).

-15-

Table 3
Sensitivity to Antifungal Compounds ^a
(Minimum Inhibitory Concentration, mg/ml)

5	Compound	YPH499 ^b	Strain A	Strain B
	Amphotericin B	2.5	2.5	2.5
	Nystatin	20	20	20
	ECB analog ^c	1.2	1.2	1.2
	Cycloheximide	0.3	0.3	0.6
10	Fluconazole	>80	>80	>80
	Itraconazole	1.2	1.2	1.2

^a Cells were grown in SC-URA (2% glucose)

^b YPH499 is transformed by YEp352.

15 ^c Echinocandin analog

-16-

SEQUENCE LISTING

- 5 (1) GENERAL INFORMATION:
- (i) APPLICANT: Heidler, Steven A.
Radding, Jeffrey A.
Young, Debra A.
- 10 (ii) TITLE OF INVENTION: Screen For Inhibitors of Fungal IPC
Synthase
- (iii) NUMBER OF SEQUENCES: 10
- 15 (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Eli Lilly and Company
(B) STREET: Lilly Corporate Center
(C) CITY: Indianapolis
20 (D) STATE: Indiana
(E) COUNTRY: U.S.
(F) ZIP: 46285
- (v) COMPUTER READABLE FORM:
25 (A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- 30 (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:
- 35 (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Webster, Thomas D.
(B) REGISTRATION NUMBER: 39872
(C) REFERENCE/DOCKET NUMBER: X-11211
- 40 (ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: 317-276-3334
- (2) INFORMATION FOR SEQ ID NO:1:
- 45 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
50 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "oligonucleotide"
- 55 (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- 60 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:
- GCAATTACGG ATCCGGTTGG TCTTATGTAG ATAC
- 65 (2) INFORMATION FOR SEQ ID NO:2:

-17-

(i) SEQUENCE CHARACTERISTICS:
5 (A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

10 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

20 GAATACGCAT AACTCGAGAG GATGATTTCT GATTAGG 37

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:
25 (A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

30 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

40 TTCAAGGATC CTTGGGCCAA AAGCTATACC 30

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:
45 (A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

50 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

60 CATGGTAATC TAGATCCTCT GAAACCTCTG C 31

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:
65 (A) LENGTH: 29 base pairs

-18-

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: other nucleic acid
(iii) HYPOTHETICAL: NO
10 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:
GCGCTCTAGATTGCCTCTGCAAAAGTTCC

15 (2) INFORMATION FOR SEQ ID NO:6:
(i) SEQUENCE CHARACTERISTICS:
20 (A) LENGTH: 41 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
25 (iii) HYPOTHETICAL: NO
(iv) ANTI-SENSE: NO

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:
CAGAGTAAAAAAGGGAAGCTTACGAAAAAAATTTCGTAAGG

(2) INFORMATION FOR SEQ ID NO:7:
35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 35 base pairs
(B) TYPE: nucleic acid
40 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
(iii) HYPOTHETICAL: NO
45 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:
50 CTTCTTAAATCAACAAGCTTTAAGAATATATTTCC

(2) INFORMATION FOR SEQ ID NO:8:
(i) SEQUENCE CHARACTERISTICS:
55 (A) LENGTH: 28 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
60 (ii) MOLECULE TYPE: other nucleic acid

-19-

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GCGCTCGAGGATTTTCGTGCAAACAAGC

10 (2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 base pairs

(B) TYPE: nucleic acid

15 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

20 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATTCTCTCGAGCAATGTCTGCTCTTTCG

30 (2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 base pairs

(B) TYPE: nucleic acid

35 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

40 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

ACACAAGGATCCACATCATTCCAATACC

-20-

WE CLAIM:

1. A biologically pure culture of yeast, said culture comprising a strain carrying a recombinantly produced deletion mutation at the *aur1*⁺ chromosomal locus
5 wherein said strain has been transformed with an expression vector comprising a vector-borne *aur1*⁺ gene that is operably-linked to a heterologous promoter such that expression of said vector-borne *aur1*⁺ gene rescues said strain from non-viability.
- 10 2. A culture, as in Claim 1 wherein said yeast is *Saccharomyces cerevisiae* or *Schizosaccharomyces pombe*.
3. A culture, as in claim 1, wherein said vector-borne *aur1*⁺ gene is operably-linked to a TPI promoter, GAL promoter, or nmt promoter.
15
4. A whole-cell screen for inhibitors of fungal IPC synthase comprising the steps of:
20
 - a. growing a suitable strain of fungal cells in a suitable medium;
 - b. exposing, by any suitable means, said cells to test compounds; and
 - c. ascertaining which of said compounds inhibits
25 the growth of said cells.
5. A whole-cell screen for inhibitors of fungal IPC synthase comprising the steps of:
30
 - a. growing in a suitable medium a first culture of fungal cells and a second culture of fungal cells wherein said first culture comprises cells that exhibit IPC synthase activity at or near the level of wild-type cells, and wherein said second culture comprises cells that exhibit altered IPC synthase activity such that said second culture
35 is hypersensitive to IPC synthase inhibitors;
 - b. exposing, by any suitable means, said cultures to test compounds; and

-21-

c. ascertaining by any suitable means the degree to which said compounds inhibit the growth of said cultures.

6. A whole-cell screen for inhibitors of fungal
5 IPC synthase comprising the steps of:
- a. growing in a suitable medium a first culture of wild-type *Saccharomyces cerevisiae* cells and a second culture comprising a culture of *Saccharomyces cerevisiae* of Claim 2 wherein said second culture is hypersensitive to IPC
10 synthase inhibitors;
 - b. exposing, by any suitable means, said cultures to test compounds; and
 - c. ascertaining by any suitable means the degree to which said compounds inhibit the growth of said cultures.

15

7. A whole-cell screen for inhibitors of fungal
IPC synthase comprising the steps of:
- a. growing in a suitable medium a first culture of wild-type *Schizosaccharomyces pombe* cells and a second
20 culture comprising a culture of *Schizosaccharomyces pombe* of Claim 2 wherein said second culture is hypersensitive to IPC synthase inhibitors;
 - b. exposing, by any suitable means, said cultures to test compounds; and
 - 25 c. ascertaining by any suitable means the degree to which said compounds inhibit the growth of said cultures.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/06897

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :C12N 15/09

US CL :435/29, 471, 477, 254.1, 254.11, 254.3

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/29, 471, 477, 254.1, 254.11, 254.3

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS and Dialog-Biotech Directory

search terms: aur1, aureobasidium, cervisiae, pombe, inositol (3w) ceramide, IPC (w) synthase

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	HEIDLER et al. "The AUR1 gene in Saccharomyces cerevisiae encodes dominant resistance to the antifungal agent Aureobasidin A (LY295337)" Antimicrobial Agents and Chemotherapy December 1995 Vol. 39, No 12, pages 2765-2769	1-7
Y	EPO 0 644 262 A2 (TAKARA SHUZO CO., LTD.) 22 March 1995 Page 9, lines 35-43, and page 58, claims 8 and 9.	1-7

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

07 JUNE 1998

Date of mailing of the international search report

11 AUG 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

BRADLEY S. MAYHEW

Telephone No. (703) 308-0196